Méningites et Encéphalites

Que faire des recommandations ?

Pr Romain Sonnevile - Dr Yousra Kherabi

BUA dans les infections communautaires graves

Vendredi 6 Juin 2025

De quoi allons-nous parler ?

- Méningite bactérienne
- Encéphalite d'allure 'virale'
- Méningite tuberculeuse
- Méningite à cryptocoque

De quoi allons-nous parler ?

- **Méningite bactérienne**
- Encéphalite d'allure 'virale'
- Méningite tuberculeuse
- Méningite à cryptocoque

Vignette clinique 1

Y

- Homme 40 ans
- ATCD: 0, alcool occasionnel
- Histoire récente : fièvre, céphalées depuis 3 jours....
- Retrouvé somnolent au domicile par sa femme
- Au SAU :
 - Température 39.1°C GCS = 13
 - Pas de syndrome méningé
 - Pas de signe neurologique focal

 - PA 130/80 mmHg, FC 110/mn
 - FR 20, SpO2 99% en air ambiant

Recommandations SPILF 2017

Grade A : Toute dysfonction aiguë du SNC associée à de la fièvre doit faire évoquer une méningo-encéphalite infectieuse

Conduite à tenir à l'arrivée: laquelle de ces propositions est juste?

- A. Vous débutez un traitement antibiotique avant toute chose
- B. Vous réalisez une ponction lombaire (PL) d'emblée
- C. Vous réalisez une imagerie cérébrale puis une PL
- D. Vous laissez le patient en surveillance simple car l'absence de syndrome méningé clinique rend le diagnostic de méningite bactérienne peu probable

Vignette clinique

- PL : pression ouverture élevée (> 20 cm H₂O)
- Traitement initial, débuté dès la constatation d'un LCS trouble:
 - Dexaméthasone 10 mg x 4 / j IVD
 - Cefotaxime 4g puis 4 g x 4 / j IVL

**** CSF Interpretation

Investigation	Normal	Bacterial	Viral	Tuberculous	Fungal
Opening pressure	10-20cm	Highly Cloudy	Normal/high	High	High/very high
Colour	Clear	Cloudy	“Gin” clear	Cloudy/yellow	Clear/cloudy
Cells	<5	High/very high 100-50000	Slightly increased 5-1000	Slightly increased <500	Normal-high 0-1000
Differential	Lymphocytes	Neutrophils	Lymphocytes	Lymphocytes	Lymphocytes
CSF/Plasma Glucose	50-66%	Low<40%	Normal	Low-very low (30%)	Normal-low
Protein (g/l)	<0.45	High >1	Normal-high 0.5-1	High-very high 1.0-5.0	Normal-high 0.2-5.0

Bactéries les plus fréquentes

ESCMID guideline: diagnosis and treatment of acute bacterial meningitis

D. van de Beek¹, C. Cabellos², O. Dzunova³, S. Esposito⁴, M. Klein⁵, A. T. Kloek¹, S. L. Leib⁶, B. Mourvillier⁷, C. Ostergaard⁸, P. Pagliano⁹, H. W. Pfister⁵, R. C. Read¹⁰, O. Resat Sipahi¹¹ and M. C. Brouwer¹, for the ESCMID Study Group for Infections of the Brain (ESGIB)

Grade A recommendation

In patients with bacterial meningitis, **classic clinical characteristics may be absent.**

Bacterial meningitis should not be ruled out solely on the absence of classic symptoms.

ESCMID guideline: diagnosis and treatment of acute bacterial meningitis

D. van de Beek¹, C. Cabellos², O. Dzunova³, S. Esposito⁴, M. Klein⁵, A. T. Kloek¹, S. L. Leib⁶, B. Mourvillier⁷, C. Ostergaard⁸, P. Pagliano⁹, H. W. Pfister⁵, R. C. Read¹⁰, O. Resat Sipahi¹¹ and M. C. Brouwer¹, for the ESCMID Study Group for Infections of the Brain (ESGIB)

Grade A recommendation

It is strongly recommended to perform **cranial imaging before LP** in patients with:

- focal neurologic deficits (excluding cranial nerve palsies)
- new-onset seizures
- severely altered mental status (GCS < 10)
- severely immunocompromised state

In patients lacking these characteristics, cranial imaging is NOT recommended.

Critical care management of infectious meningitis and encephalitis

Geert Meyfroidt^{1,2*}, Pedro Kurtz³ and Romain Sonnevile^{4,5}

ESCMID guideline: diagnosis and treatment of acute bacterial meningitis

D. van de Beek¹, C. Cabellos², O. Dzupova³, S. Esposito⁴, M. Klein⁵, A. T. Kloek¹, S. L. Leib⁶, B. Mourvillier⁷, C. Ostergaard⁸, P. Pagliano⁹, H. W. Pfister⁵, R. C. Read¹⁰, O. Resat Sipahi¹¹ and M. C. Brouwer¹, for the ESCMID Study Group for Infections of the Brain (ESGIB)

Grade A recommendation

It is strongly recommended to start antibiotic therapy **as soon as possible** in acute bacterial meningitis patients.

The time period until antibiotics are **administered should not exceed 1 hour**.

Whenever lumbar puncture is delayed (e.g. cranial CT), **empirical treatment should be started immediately** based on clinical suspicion, even if the diagnosis has not been established.

Antibiothérapies probabilistes pour les méningites présumées bactériennes

- Cefotaxime : 200 mg/kg/day IV (intermittent infusion, 4 doses) for all patients
- In countries where the prevalence of cephalosporin-R pneumococcus is > 1%, add :
 - Vancomycin : 30 mg/kg IV bolus followed by continuous infusion 45-60 mg/kg/day
 - or
 - Rifampicin : 600 mg / day IV (intermittent infusion)
- if risk factor(s) for *Listeria monocytogenes* is present, add :
(Age > 50 years, alcoholism, cancer, immunosuppressive therapy...)
 - Amoxicillin : 200 mg/kg/day IV (intermittent infusion, 4 doses)

Clinical Features and Prognostic Factors in Adults with Bacterial Meningitis

Diederik van de Beek, M.D., Ph.D., Jan de Gans, M.D., Ph.D.,
Lodewijk Spanjaard, M.D., Ph.D., Martijn Weisfelt, M.D.,
Johannes B. Reitsma, M.D., Ph.D., and Marinus Vermeulen, M.D., Ph.D.

- **Gram's staining of cerebrospinal fluid** revealed the microorganism in 524 of 652 episodes (sensitivity, 80 percent; specificity, 97 percent).
- The yield of **Gram's staining** was similar in patients who had previously received antimicrobial therapy and those who had not.
- **Positive blood cultures** were observed in **404 / 611 (66%) patients**

A propos des corticoïdes: laquelle (ou lesquelles) de ces propositions est (sont) juste(s)?

- A. Ils sont indiqués dans les méningites bactériennes
- B. Ils réduisent la mortalité des méningites à méningocoque
- C. Ils réduisent la mortalité des méningites à pneumocoque
- D. Ils sont associés à une réduction des séquelles neurosensorielles
- E. La durée de traitement standard chez l'adulte est de 8 jours

DEXAMETHASONE IN ADULTS WITH BACTERIAL MENINGITIS

JAN DE GANS, PH.D., AND DIEDERIK VAN DE BEEK, M.D., FOR THE EUROPEAN DEXAMETHASONE IN ADULTHOOD BACTERIAL MENINGITIS STUDY INVESTIGATORS*

OUTCOME AND CULTURE RESULTS	DEXAMETHASONE GROUP	PLACEBO GROUP	RELATIVE RISK (95% CI)†	P VALUE
	no./total no. (%)			
Unfavorable outcome				
All patients	23/157 (15)	36/144 (25)	0.59 (0.37–0.94)	0.03
<i>Streptococcus pneumoniae</i>	15/58 (26)	26/50 (52)	0.50 (0.30–0.83)	0.006
<i>Neisseria meningitidis</i>	4/50 (8)	5/47 (11)	0.75 (0.21–2.63)	0.74
Other bacteria	2/12 (17)	1/17 (6)	2.83 (0.29–27.8)	0.55
Negative bacterial culture‡	2/37 (5)	4/30 (13)	0.41 (0.08–2.06)	0.40
Death				
All patients	11/157 (7)	21/144 (15)	0.48 (0.24–0.96)	0.04
<i>S. pneumoniae</i>	8/58 (14)	17/50 (34)	0.41 (0.19–0.86)	0.02
<i>N. meningitidis</i>	2/50 (4)	1/47 (2)	1.88 (0.76–20.1)	1.00
Other bacteria	1/12 (8)	1/17 (6)	1.42 (0.10–20.5)	1.00
Negative bacterial culture	0/37	2/30 (7)	—	0.20

Corticosteroids for acute bacterial meningitis

	RR	95% CI
Mortality*	0.90	0.80-1.01
Mortality in adults**	0.74	0.53-1.05
Mortality in patients with <i>S. pneumoniae</i> meningitis	0.84	0.72-0.98
Hearing loss	0.74	0.63-0.87
Neurological sequelae	0.83	0.69-1.00

* p= 0.07, ** p= 0.09

There was no beneficial effect of corticosteroid therapy in low-income countries.

ESCMID guideline: diagnosis and treatment of acute bacterial meningitis

D. van de Beek¹, C. Cabellos², O. Dzunova³, S. Esposito⁴, M. Klein⁵, A. T. Kloek¹, S. L. Leib⁶, B. Mourvillier⁷, C. Ostergaard⁸, P. Pagliano⁹, H. W. Pfister⁵, R. C. Read¹⁰, O. Resat Sipahi¹¹ and M. C. Brouwer¹, for the ESCMID Study Group for Infections of the Brain (ESGIB)

Grade A recommendations

Empiric treatment with dexamethasone is strongly recommended for all adults

(10 mg q6h for 4 days) with acute bacterial meningitis in the setting of high income countries

Treatment with dexamethasone is **strongly recommended to be initiated with the first dose of antibiotic treatment.**

Steroid use in non-pneumococcal and non-*Haemophilus* bacterial meningitis – Authors' reply

To conclude, dexamethasone should be initiated with the first dose of antibiotics in all patients with community-acquired bacterial meningitis beyond the neonatal age. On the basis of available evidence, we advise to continue dexamethasone treatment in this patient group for 4 days regardless of microbial cause, except in patients with *Listeria monocytogenes*.

Méningite - take home messages

- Au SAU / en réa souvent, pas la séquence idéale PL puis ATB
L'URGENCE est à l'antibiothérapie < 1 heure
- Dexaméthasone avec la première dose d'antibiotique
En 2025, pour toute méningite bactérienne... sauf *Listeria*

De quoi allons-nous parler ?

- Méningite bactérienne
- **Encéphalite d'allure 'virale'**
- Méningite tuberculeuse
- Méningite à cryptocoque

Vignette clinique 2

H 27 ans, confusion, Fièvre 38.7° C

PL : 80 éléments / mm³, 90% lymphocytes, prot 0.8g/l, glycorachie 4 mmol/l

Quelle est votre attitude thérapeutique ?

**** CSF Interpretation

Investigation	Normal	Bacterial	Viral	Tuberculous	Fungal
Opening pressure	10-20cm	Highly Cloudy	Normal/high	High	High/very high
Colour	Clear	Cloudy	“Gin” clear	Cloudy/yellow	Clear/cloudy
Cells	<5	High/very high 100-50000	Slightly increased 5-1000	Slightly increased <500	Normal-high 0-1000
Differential	Lymphocytes	Neutrophils	Lymphocytes	Lymphocytes	Lymphocytes
CSF/Plasma Glucose	50-66%	Low<40%	Normal	Low-very low (30%)	Normal-low
Protein (g/l)	<0.45	High >1	Normal-high 0.5-1	High-very high 1.0-5.0	Normal-high 0.2-5.0

Recommandations SPILF 2017

Grade A : en l'absence d'orientation étiologique, le traitement initial associe :

Aciclovir : 10mg/kg/8h IVL (dans les 6h suivant admission)

ET

Amoxicilline : 200 mg/kg/24h IVL en perfusion continue

Adapter les posologies à la fonction rénale

Réévaluer à 48 heures +++

(cultures bactériennes, PCR HSV, VZV, enterovirus)

Recommandations SPILF 2017

Grade A : si elle est possible, l'IRM cérébrale est l'imagerie à réaliser en première intention (Diffusion, FLAIR, T2*, T1 gadolinium)

H 27 ans, confusion, Fièvre 38.7° C

PL : 80 éléments / mm³, 90% lymphocytes, prot 0.8g/l, glycorachie 4 mmol/l

Aciclovir IV 10mg/kg/8h IVL

DEXENCEPH HSV Encephalitis DXM 10 mg x 4 / j versus placebo : no effect on cognitive function

	Dexamethasone		Control		Adjusted mean difference in Dexamethasone vs Control (95% CI)	P-value
	Mean	N	Mean	N		
Primary outcome						
WMS-IV Auditory Memory Index (Primary Outcome)	71 (26)	42	69 (25)	39	1.7 (-10, 13)	.756
Secondary outcomes						
WMS-IV Visual Memory Index (VMI)	69 (25)	33	71 (26)	31	-1.1 (-13, 11)	.861
WMS-IV Immediate Memory Index (IMI)	71 (25)	33	67 (24)	31	-1.2 (-14, 12)	.853
WMS-IV Delayed Memory Index (DMI)	68 (24)	33	64 (24)	31	-0.1 (-13, 13)	.997
WAIS-IV Working Memory Index (WMI)	85 (25)	36	82 (27)	36	4.6 (-8, 17)	.465
WAIS-IV Processing Speed Index (PSI)	80 (23)	33	77 (26)	32	4 (-8, 16)	.503
Trail Making Test A	96 (99)	34	134 (119)	35	-38 (-89, 13)	.138
Trail Making Test B	165 (96)	33	202 (106)	34	-33.7 (-80, 13)	.153
Perceived deficits questionnaire (PDQ)	30 (17)	29	28 (14)	25	2.5 (-6, 11)	.574
Beck Anxiety Inventory (BAI)	14 (11)	27	11 (11)	26	3.3 (-3, 10)	.295
Beck Depression Inventory (BDI)	17 (12)	27	14 (11)	26	2.7 (-4, 9)	.408

Dans quelques jours ...

EAN-ESCMID guidelines on the diagnosis and management of encephalitis in adults caused by infection

VZV : 15 mg/kg/8h → 10 mg/kg/8h

Encéphalite virale - take home messages

- Importance de l'imagerie ++
- CTC et HSV : on attend les résultats définitifs de DEXENCEPH
- Les nouvelles recommandations ESCMID arrivent : baisse de la posologie pour VZV, pas de nouveauté sur la CTC

De quoi allons-nous parler ?

- Méningite bactérienne
- Encéphalite d'allure 'virale'
- **Méningite tuberculeuse**
- Méningite à cryptocoque

Vignette clinique 3

Madame M. 49 ans consulte aux Urgences pour céphalées intenses depuis 8 jours

ATCD :

Infection par le VIH en rupture de suivi

Dernier bilan immuno-virologique 11/2024 : 80 CD4 CV 5 log

Cliniquement :

T°38.7 GCS 14

PA 113/71 mmHg - Fc 94 bpm - FR 23/min - SpO2 96% en AA

PL : 57 éléments (91% lymphocytes), 0 hématies, protéinorachie 1.8 g/L, glycorachie 1,9 mmol/L (glycémie : 4,9 mmol/L)

Leptoméningite base du crâne

Plusieurs lésions nodulaires cérébelleuses

Miliaire tuberculeuse pulmonaire : BK crachat + à 1 BAAR/lame

Quel traitement instaurez-vous ?

- A) Quadrithérapie RMP INH PZA EMB per os posologie standard
- B) Quadrithérapie RMP INH PZA EMB IV posologie standard
- C) Quadrithérapie RMP INH PZA EMB per os posologie élevée RMP
- D) Quadrithérapie RMP INH PZA EMB per os posologie élevée INH
- E) Corticothérapie systémique

**** CSF Interpretation

Investigation	Normal	Bacterial	Viral	Tuberculous	Fungal
Opening pressure	10-20cm	Highly Cloudy	Normal/high	High	High/very high
Colour	Clear	Cloudy	“Gin” clear	Cloudy/yellow	Clear/cloudy
Cells	<5	High/very high 100-50000	Slightly increased 5-1000	Slightly increased <500	Normal-high 0-1000
Differential	Lymphocytes	Neutrophils	Lymphocytes	Lymphocytes	Lymphocytes
CSF/Plasma Glucose	50-66%	Low<40%	Normal	Low-very low (30%)	Normal-low
Protein (g/l)	<0.45	High >1	Normal-high 0.5-1	High-very high 1.0-5.0	Normal-high 0.2-5.0

TUER LE BK

Grade A : il est recommandé d'instaurer un traitement d'épreuve anti-tuberculeux si le tableau clinique, biologique et l'imagerie sont à ce stade en faveur d'un diagnostic de tuberculose, même en cas d'examen microscopique et de PCR négatifs.

Une histoire
différente en
soins intensifs :

Les molécules
ne se valent pas
toutes

Ttt antituberculeux initial:

- ✓ Jusqu'aux résultats de l'antibiogramme
 - Isoniazide (I) (5mg/kg/j)
 - Rifampicine (R) (10 mg/kg/j)
 - Pyrazinamide (P) (30 mg/kg/j sans dépasser 2 g)
 - Ethambutol (E) (20 mg/kg/j)
- ✓ Arrêt de l'éthambutol si souche sensible à l'isoniazide
- ✓ Durée initiale de 2 mois
- ✓ Puis **bithérapie par IR pour une durée totale de 12 mois**

BHE : tous les anti-BK ne se valent pas

Ruslami et al LID 2013

	Standard daily dose for adults	Estimated ratio of CSF to plasma concentration	Comments
Isoniazid	300 mg	80–90%	Essential drug; good CSF penetration throughout treatment
Rifampicin	450 mg (weight <50 kg) or 600 mg (weight ≥50 kg)	10–20%	Essential drug, despite relatively poor CSF penetration; higher doses might improve effectiveness
Pyrazinamide	1.5 g (weight <50 kg) or 2.0 g (weight ≥50 kg)	90–100%	Excellent CSF penetration throughout treatment
Ethambutol	15 mg/kg	20–30%	Poor CSF penetration once meningeal inflammation resolves
Streptomycin	15 mg/kg (1 g maximum)	10–20%	Poor CSF penetration once meningeal inflammation resolves
Kanamycin	15 mg/kg	10–20%	Poor CSF penetration once meningeal inflammation resolves
Amikacin	15–20 mg/kg	10–20%	Poor CSF penetration once meningeal inflammation resolves
Moxifloxacin	400 mg	70–80%	Good CSF penetration
Levofloxacin	1000 mg	70–80%	Good CSF penetration
p-Aminosalicylic acid	10–12 g	No data	Probably very poor CSF penetration unless meninges are inflamed
Ethionamide or protionamide	15–20 mg/kg (1 g maximum)	80–90%	Good CSF penetration
Cycloserine	10–15 mg/kg	80–90%	Good CSF penetration
Linezolid	1200 mg	40–70%	Variable interindividual CSF pharmacokinetics
Capreomycin	15–20 mg/kg	No data	..

Table 4: CSF penetration of first-line and second-line antituberculosis drugs^{82–85}

Augmenter la dose d'INH : pourquoi pas ?

ORIGINAL ARTICLE

Intensified Antituberculosis Therapy in Adults with Tuberculous Meningitis

A. Dorothee Heemskerck, M.D., Nguyen D. Bang, Ph.D., Nguyen T.H. Mai, Ph.D., Tran T.H. Chau, Ph.D., Nguyen H. Phu, Ph.D., Pham P. Loc, M.D., Nguyen V.V. Chau, Ph.D., Tran T. Hien, Ph.D., Nguyen H. Dung, Ph.D., Nguyen T.N. Lan, Ph.D., Nguyen H. Lan, M.D., Nguyen N. Lan, M.D., Le T. Phong, M.D., Nguyen N. Vien, M.D., Nguyen Q. Hien, M.D., Nguyen T.B. Yen, M.D., Dang T.M. Ha, Ph.D., Jeremy N. Day, F.R.C.P., Maxine Caws, Ph.D., Laura Merson, B.S., Tran T.V. Thinh, M.D., Marcel Wolbers, Ph.D., Guy E. Thwaites, F.R.C.P., and Jeremy J. Farrar, F.R.C.P.

60 days: Rifampicin 15mg/kg/day + Levofloxacin + HZE, then 10HZ

Pharmacokinetics and Pharmacodynamics of Intensive Antituberculosis Treatment of Tuberculous Meningitis

Junjie Ding^{1,2,3}, Nguyen Thuy Thuong Thuong⁴, Toi Van Pham⁴, Dorothee Heemskerck⁴, Thomas Pouplin^{1,3}, Chau Thi Hong Tran⁵, Mai Thi Hoang Nguyen⁵, Phu Hoan Nguyen^{4,5}, Loc Phu Phan⁵, Chau Van Vinh Nguyen⁵, Guy Thwaites^{1,4} and Joel Tarning^{1,2,3*}

Isoniazid exposure associated with outcomes; not rifampicin exposure

N Engl J Med. 2016 Jan 14;374(2):124-34
Clin Pharmacol Ther. 2020 Apr;107(4):1023-1033.

Augmenter la dose de RMP : plutôt non

Wellcome Open Research Wellcome Open Research 2020, 4:190 Last updated: 13 NOV 2024

STUDY PROTOCOL

REVISED **High dose oral rifampicin to improve survival from adult tuberculous meningitis: A randomised placebo-controlled double-blinded phase III trial (the HARVEST study)**

[version 2; peer review: 2 approved]

Suzaan Marais ^{1*}, Fiona V Cresswell ^{2-4*}, Raph L. Hamers ^{5,6}, Lindsey H.M. te Brake⁷, Ahmad R. Ganiem^{8,9}, Darma Imran ¹⁰, Ananta S. Bangdiwala ¹¹, Emily Martyn², John Kasibante ², Enock Kagimu ², Abdu K. Musubire ², Kartika Maharani¹⁰, Riwanti Estiasari¹⁰, Ardiana Kusumaningrum ¹², Nadytia Kusumadjayanti⁹, Vycke Yunivita^{8,9}, Kogieleum Naidoo^{13,14}, Richard R. Lessells ^{13,15}, Yunus Moosa¹⁶, Elin M. Svensson ^{7,17}, Katherine Huppler Hullsiek ¹¹, Rob E. Aarnoutse⁷, David R. Boulware ¹⁸, Reinout van Crevel ^{6,19}, Rovina Ruslami^{9,20*}, David B. Meya ^{2*}

Late Breakers

Trials in severe infections

CLN

Download calendar even

Chair(s)

Lennie DERDE, Netherlands

Chair

Steven TONG, Australia

Chair

Last talk in the session!

High-dose rifampicin in the treatment of tuberculous meningitis: results of the HARVEST phase III multi-country randomised clinical trial

Reinout VAN CREVEL, Netherlands

Prendre en compte plusieurs dimensions

Advancing the chemotherapy of tuberculous meningitis: a consensus view

Pondérer le choix des antibiotiques par un score combiné associant :

- Activité bactéricide
- Efficacité clinique prouvée dans essais contrôlés
- Concentration dans le LCS
- Tolérance
- Interaction médicamenteuse
- Facilité d'accès (prix, voie d'administration, réglementation etc.)

	Anti-tuberculosis activity	Clinical efficacy in tuberculous meningitis	Site of disease exposure	Safety and tolerability	Drug-drug interactions	Access
Rifampicin	3	2	3	2	0	2
Isoniazid	3	2	3	2	1	2
Linezolid ³¹	2	2	3	1	1	2
Fluoroquinolones ³²	3	2	3	2	1	2
Pyrazinamide	2	2	3	2	1	2
Pretomanid or delamanid ³³	3	1	3	2	1	2
Bedaquiline	3	1	2	2	0	2
Alpibectir or ethionamide	2	0	3	2	1	..
Rifabutin	3	0	2	2	0	2
Clofazimine	1	0	1	2	1	2
Ethionamide	1	2	2	NA	1	2
Cycloserine	1	0	2	NA	1	2
Ethambutol	1	NA	NA	2	1	2
DprE1 inhibitors	3	0	0	2	1	NA

Scores derived from the scoring system in table 2. NA=not applicable.

Table 3: Characteristics of individual drugs for use in tuberculous meningitis

High-dose rifampicin

- 1. RFqZL
- 2. RFqZAlpE

Rifamycin-free

- 3. BPamZ (SimpliciTB regimen)
- 4. BPaLM (TB-PRACTECAL)
- 5. BPaOS (tested in pan-tuberculosis trial)

Bedaquiline plus rifabutin

- 6. BZMRb (superior to RHZE in mice)
 - 7. BPaORb
 - 8. BPaAlpERb
- } (Modified BPaOS regimen)

Drug scores

 ≥12
 9-11
 <9
 Not currently manufactured

		Rifamycin	Isoniazid	Pyrazinamide	Fluoroquinolone	Bedaquiline	Oxazolidinone	Nitroimidazole	AlpE	DprE1
1	RFqZL	R					L			
2	RFqZAlpE	R								
3	BPamZ							Pa		
4	BPaLM						L	Pa		
5	BPaOS						S	Pa		
6	BZMRb	Rb								
7	BPaORb	Rb						Pa		
8	BPaAlpERb	Rb						Pa		

REDUIRE L'INFLAMMATION

Dexamethasone for the Treatment of Tuberculous Meningitis in Adolescents and Adults

The NEW ENGLAND JOURNAL of MEDICINE

Table 3. Outcomes of 545 Patients Nine Months after Randomization.

Group	No. of Patients	Outcome			
		Good	Inter-mediate	Severe Disability	Death
Dexamethasone*	274	104 (38.0)	49 (17.9)	34 (12.4)	87 (31.8)
Placebo	271	95 (35.1)	42 (15.5)	22 (8.1)	112 (41.3)

Dexamethasone for the Treatment of Tuberculous Meningitis in Adolescents and Adults

Outcome and Group	Dexamethasone <i>no./total no. (%)</i>	Placebo <i>no./total no. (%)</i>	Relative Risk (95% CI)	P Value
Death				
All patients	87/274 (31.8)	112/271 (41.3)	0.69 (0.52–0.92)	0.01
Grade				
I	15/90 (16.7)	26/86 (30.2)	0.47 (0.25–0.90)	0.02
II	38/122 (31.1)	50/125 (40.0)	0.71 (0.46–1.1)	0.11
III	34/62 (54.8)	36/60 (60.0)	0.81 (0.51–1.29)	0.38
Relative risk of death stratified according to grade†			0.68 (0.52–0.91)	0.007
HIV status				
Negative	57/227 (25.1)	67/209 (32.1)	0.72 (0.51–1.02)	0.07
Positive	27/44 (61.4)	37/54 (68.5)	0.86 (0.52–1.41)	0.55
Undetermined	3/3 (100)	8/8 (100)	1.16 (0.71–1.91)	0.71
Relative risk of death stratified according to HIV status‡			0.78 (0.59–1.04)	0.08

Quel protocole ?

WHO ^{79,80} and UK ⁵² recommendations				
	Daily dose in children	Daily dose in adults	Route of administration	Duration
Adjunctive corticosteroids				
Prednisolone	4 mg/kg*	2.5 mg/kg*	Intravenous initially, then switch to oral when safe to do so	4 weeks then reduce to stop over 4 weeks
Dexamethasone	0.6 mg/kg*	0.4 mg/kg*	Intravenous initially, then switch to oral when safe to do so	Reducing each week to stop over 6–8 weeks

Adjunctive Dexamethasone for Tuberculous Meningitis
in HIV-Positive Adults

Joseph Donovan, Ph.D., Nguyen D. Bang, Ph.D., Darma Imran, M.D., Ho D.T. Nghia, Ph.D., Erlina Burhan, Ph.D.,
Dau T.T. Huong, M.Sc., Nguyen T.T. Hiep, M.D., Lam H.B. Ngoc, B.Sc., Dang V. Thanh, M.D.,
Nguyen T. Thanh, M.D., Anna L.S. Wardhani, B.Sc., Kartika Maharani, M.D., Cakra P. Gasmara, M.D.,
Nguyen H.H. Hanh, M.D., Pham K.N. Oanh, M.D., Riwanti Estiasari, Ph.D., Do D.A. Thu, B.Sc.,
Ardiana Kusumaningrum, M.D., Le T. Dung, M.D., Do C. Giang, Ph.D., Dang T.M. Ha, Ph.D.,
Nguyen H. Lan, M.D., Nguyen V.V. Chau, Ph.D., Nguyen T.M. Nguyet, B.Sc., Ronald B. Geskus, Ph.D.,
Nguyen T.T. Thuong, Ph.D., Evelyne Kestelyn, M.P.H., Raph L. Hamers, Ph.D., Nguyen H. Phu, Ph.D.,
and Guy E. Thwaites, F.R.C.P., for the ACT HIV Investigators*

Table 1. Baseline Characteristics in the Intention-to-Treat Population.*

Characteristic	Total (N=520)	Dexamethasone (N=263)	Placebo (N=257)
Median age (IQR) — yr	36 (30–41)	36 (29–41)	36 (30–42)
Male sex — no. (%)	396 (76.2)	208 (79.1)	188 (73.2)
Diagnostic category — no. (%)†			
Definite tuberculous meningitis	212 (40.8)	108 (41.1)	104 (40.5)
Probable tuberculous meningitis	253 (48.7)	129 (49.0)	124 (48.2)
Possible tuberculous meningitis	52 (10.0)	24 (9.1)	28 (10.9)
Not tuberculous meningitis‡	0	0	0
Unknown§	3 (0.6)	2 (0.8)	1 (0.4)
Modified MRC disease-severity grade — no. (%)¶			
I	196 (37.7)	99 (37.6)	97 (37.7)
II	251 (48.3)	125 (47.5)	126 (49.0)
III	73 (14.0)	39 (14.8)	34 (13.2)
Median score on the Glasgow Coma Scale (IQR)	14 (12–15)	14 (12–15)	14 (12–15)
CSF microbiologic tests — no./total no. (%)			
Positive Ziehl–Neelsen stain	100/504 (19.8)	52/255 (20.4)	48/249 (19.3)
Positive Xpert MTB/RIF test	107/401 (26.7)	53/204 (26.0)	54/197 (27.4)
Positive Xpert MTB/RIF Ultra test**	46/113 (40.7)	22/56 (39.3)	24/57 (42.1)
Positive mycobacterial culture	148/508 (29.1)	77/256 (30.1)	71/252 (28.2)
ART status at enrollment — no. (%)			
No previous ART	255 (49.0)	133 (50.6)	122 (47.5)
ART for >3 mo	104 (20.0)	46 (17.5)	58 (22.6)
ART of undetermined duration††	106 (20.4)	58 (22.1)	48 (18.7)
Unknown status or missing data	55 (10.6)	26 (9.9)	29 (11.3)
CD4 cell count at enrollment — no./total no. (%)			
≤50 per cubic millimeter	251/484 (51.9)	126/244 (51.6)	125/240 (52.1)
51 to 100 per cubic millimeter	89/484 (18.4)	45/244 (18.4)	44/240 (18.3)
101 to 200 per cubic millimeter	71/484 (14.7)	36/244 (14.8)	35/240 (14.6)
>200 per cubic millimeter	73/484 (15.1)	37/244 (15.2)	36/240 (15.0)

A Death from Any Cause, Intention-to-Treat Population

No. at Risk

Dexamethasone	263	202	182	172	166	161	156	154	151	149	146	143	139
Placebo	257	185	165	152	147	141	137	135	134	134	133	130	127

B Between-Group Differences in Survival, Intention-to-Treat Population

Figure 2. Death from Any Cause during 12 Months after Randomization.

Table 2. Death from Any Cause in Prespecified Subgroups in the Intention-to-Treat Population.^{a,b}

Subgroup	Dexamethasone (N = 263)	Placebo (N = 257)	Hazard Ratio (95% CI)
	<i>no. of deaths/no. of participants</i>		
Overall	116/263	126/257	0.85 (0.66–1.10) [†]
Modified MRC disease-severity grade			
I	22/99	28/97	0.72 (0.41–1.25)
II	60/125	68/126	0.82 (0.58–1.16)
III	34/39	30/34	1.03 (0.63–1.69)
Diagnostic category			
Definite tuberculous meningitis	48/108	49/104	0.90 (0.61–1.35)
Probable tuberculous meningitis	61/129	61/124	0.91 (0.64–1.30)
Possible tuberculous meningitis	5/24	15/28	0.34 (0.12–0.94)
LTA4H genotype [‡]			
TT	12/25	11/26	1.06 (0.47–2.41)
CT	49/117	59/114	0.72 (0.49–1.05)
CC	38/84	36/80	1.04 (0.66–1.63)
Antituberculosis-drug resistance [§]			
MDR or rifampin mono-resistance	7/10	5/6	0.66 (0.21–2.11)
Isoniazid resistance without MDR	6/14	13/20	0.56 (0.21–1.49)
No or other resistance	22/52	13/45	1.58 (0.79–3.13)
ART status at enrollment			
No previous ART	64/133	64/122	0.85 (0.60–1.21)
>3 Mo of ART	20/46	26/58	0.96 (0.54–1.72)
ART of undetermined duration	19/58	19/48	0.80 (0.42–1.51)
CD4 cell count at enrollment			
≤50 per cubic millimeter	67/126	67/125	0.96 (0.69–1.35)
51–100 per cubic millimeter	12/45	19/44	0.52 (0.25–1.06)
101–200 per cubic millimeter	14/36	13/35	1.04 (0.49–2.22)
>200 per cubic millimeter	11/37	15/36	0.70 (0.32–1.52)

Aspirine ou clopidogrel : plutôt non

A Randomised Trial to Assess the Efficacy of Add on Therapy with Aspirin or Clopidogrel to the Standard Medical Therapy Alone in Patients with Tubercular Meningitis: ACT TBM

26 Pages • Posted: 18 Dec 2024

Interpretation: No difference was observed by an add-on therapy of antiplatelets to the standard care compared to standard care alone on imaging-based cerebral infarction, clinical stroke, or mortality.

Méningite TB - take home messages

- TBM : 50 % de mortalité dans le monde, 100 cas par an en France
- Performances insuffisantes des outils diagnostiques actuels : combiner les tests + gros volumes de LCS
- IREP - durée 12 mois
Doses, administration, combinaisons : incertains, essais en cours
- CTC : Réduction de la mortalité chez le non VIH seulement
En cours d'évaluation : aspirine, anti TNF alpha
- Nouvelles recommandations SPILF 2025 à surveiller

De quoi allons-nous parler ?

- Méningite bactérienne
- Encéphalite d'allure 'virale'
- Méningite tuberculeuse
- **Méningite à cryptocoque**

Vignette clinique 4

Madame C. 42 ans consulte aux Urgences pour céphalées intenses depuis 8 jours

ATCD :

Infection par le VIH en rupture de suivi

Bilan immuno-virologique fait en ville par médecin généraliste 05/2025 : 0 CD4 CV 6 log

Cliniquement :

T°39 GCS 14

PA 113/71 mmHg - Fc 94 bpm - FR 23/min - SpO2 96% en AA

PL : 177 éléments (93% lymphocytes), 0 hématies, protéinorachie 2.8 g/L, glycorachie 2.4 mmol/L (glycémie : 4,1 mmol/L)

Vignette clinique 4

Appel du laboratoire :

Vignette clinique 4

R

Quel élément pronostic majeur vous manque-t-il quant aux résultats de la PL ?

Rappel des informations données:

PL : 177 éléments (93% lymphocytes), 0 hématies, protéinorachie 2.8 g/L, glycorachie 2.4 mmol/L (glycémie : 4,1 mmol/L)

**** CSF Interpretation

Investigation	Normal	Bacterial	Viral	Tuberculous	Fungal
Opening pressure	10-20cm	Highly Cloudy	Normal/high	High	High/very high
Colour	Clear	Cloudy	“Gin” clear	Cloudy/yellow	Clear/cloudy
Cells	<5	High/very high 100-50000	Slightly increased 5-1000	Slightly increased <500	Normal-high 0-1000
Differential	Lymphocytes	Neutrophils	Lymphocytes	Lymphocytes	Lymphocytes
CSF/Plasma Glucose	50-66%	Low<40%	Normal	Low-very low (30%)	Normal-low
Protein (g/l)	<0.45	High >1	Normal-high 0.5-1	High-very high 1.0-5.0	Normal-high 0.2-5.0

Méningite à cryptocoque en 2025

- FDR principal : déficit en immunité CELLULAIRE (HIV +++)
- 1ere cause de méningite de l'adulte en Afrique
- 20% des morts chez les PVVIH (Rajasingham LID 2022)
- Mortalité = 40% à S10
- Handicap = 70 % (Molloy NEJM 2018)

Cryptococcal Disease in Diverse Hosts

David B. Meya, M.B., Ch.B., Ph.D., and Peter R. Williamson, M.D., Ph.D.

Table 1. Performance Characteristics of Cryptococcal Diagnostic Assays in Cerebrospinal Fluid (CSF) from Persons with Suspected Meningitis.*

Diagnostic Test	Sensitivity	Specificity	Positive Predictive Value	Negative Predictive Value
			<i>percent</i>	
Cryptococcal antigen lateral flow immunochromatographic assay	99.3	99.1	99.5	98.7
CSF culture†	90.0	100	100	85.3
100 mm ³	94.2	100	100	91.2
10 mm ³	82.4	100	100	75.8
India ink microscopy	86.1	97.3	98.2	80.2
Cryptococcal antigen latex agglutination assay				
Meridian	97.8	85.9	92.6	95.5
IMMY	97.0	100	100	95.3
Metagenomic next-generation sequencing	93.5	96.0	87.8	98.0
PCR assay‡	82.0	98.0	98.0	79.0

Pression intra-crânienne

Global guideline for the diagnosis and management of cryptococcosis: an initiative of the ECMM and ISHAM in cooperation with the ASM

Mesurer la **pression d'ouverture** à chaque ponction lombaire.

Si normale: refaire **ponction la première semaine (à J2/J3 ou à J7)**

Si ≥ 20 cmH₂O : **ponctions quotidiennes (20–30 mL) pour $\downarrow \leq 20$ cmH₂O ou 50 %.**

Si persistante : envisager **dérivation chirurgicale**

Adjunctive Dexamethasone in HIV-Associated Cryptococcal Meningitis

A All Patients

No. at Risk

Placebo	226	179	143	132	125	122	116	112
Dexamethasone	224	185	146	120	109	103	98	92

Single-Dose Liposomal Amphotericin B Treatment for Cryptococcal Meningitis

Y

Jarvis JN et al. DOI: 10.1056/NEJMoa2111904

5FC, flucytosine ; CrAg, antigène cryptococcique ; FCZ, fluconazole ; L-AmB, amphotéricine B liposomale

Global guideline for the diagnosis and management of cryptococcosis: an initiative of the ECMM and ISHAM in cooperation with the ASM

Méningite crypto - take home messages

- HIV mais pas que (transplanté d'organe solide ++)
- Attention à la pression intracrânienne
- Régime single dose Ampho B maintenant recommandé
- Attention aux effets indésirables (rénaux réduits avec ampho B SD)
- Hors soin d'urgence : traitement long
consolidation et entretien

RCP neuro-infectiologique depuis 2020

- RCP mensuelle le 3^e vendredi du mois
LA PROCHAINE LE 20 JUIN 2025
- En présentiel à Bichat et en distanciel
- Neurologues (Delafontaine, Bichat),
neuro-radiologues (FOR, Bichat),
réanimateurs, infectiologues... et vous !
- Cas d'infection avérée ou suspectée du SNC

Rejoignez la mailing list
en flashant ce QR code

MERCI DE VOTRE ATTENTION

Méningite à *Listeria* et corticostéroïdes

- Effet plutôt délétère : étude de cohorte prospective française MONALISA (Charlier et al. The Lancet 2017)

252 neurolistérioses, mortalité à 48% chez ceux recevant de la DXM vs 27%

- Bénéfice : étude de cohorte prospective néerlandaise (Brouwer et al. The Lancet eClinicalMedicine 2023) --> 162 neurolistérioses, effet favorable sur le Glasgow Outcome scale ET sur mortalité

TBM et PL déplétive

Hydrocéphalie : complique jusqu'à 86 % des TBM (Chiang LID 2014)

Le plus souvent communicante

Peu de littérature sur intérêt PL déplétive

Dans la crypto : argument aussi de la baisse de la charge fongique
≠ TBM

Quelques équipes l'ont fait : Neuro CH de St Denis (Lampros et al. ID now 2025)

Cryptococcose chez le transplanté :

Tardif (mois/années après la SOT)

Les plus à risque : coeur et intestin grêle

Pas de RCT avec les SOT spécifiquement : extrapolé depuis le VIH

The use of lipid-formulations in SOT recipients with CNS cryptococcosis was independently associated with reduced mortality compared with amphotericin B.

Cryptococcus gattii*: comparison with *C. neoformans

- Retrospective cohort, 5 centers, 1995-2013
- *C. neoformans* (452); *C. gattii* (217); 61.4% male
- *C. gattii* ($p < 0.05$):
 - Time for diagnosis 52.2 vs. 36.01 d
 - No underlying disease 40.5 vs. 10.2%
 - Pulmonary involvement 60.7 vs. 32.1%
 - Female and age 52.2 vs. 36 yrs
 - Less CNS involvement and fungemia
 - Survival at M12
 - IRIS

Panel 13: Recommendations for *C gattii*

In *C gattii* CNS disease:

- (AIII) Treat the same as *C neoformans* CNS infection
- (BIII) In non-HIV patients, consider extending induction therapy to 4-6 weeks
- (AIII) Early CSF shunting is indicated for obstructive chronic hydrocephalus

Treatment of *C gattii* lung disease is summarised in the appendix (p 17).

Baddley, CID 2021

Cryptococcal IRIS

- New clinical symptoms consistent with inflammatory process
- ± during immunological and/or virological response following initiation of HAART
- ± negative microbial cultures
- No new opportunistic infections (TB), HIV-related neoplasia or drugs-related disorders (limited diagnostic capabilities in some countries)
- Increasingly reported during various non-fungal opportunistic infections
- Rarely reported during *Pneumocystis jirovecii*, *Histoplasma capsulatum*, *Coccidioides*, *Sporothrix* and *Talaromyces marneffeii* infections

*Wislez Am J Respir Crit Care Med 2001; Jenny-Avital CID 2002;
Shelburne AIDS Rev 2003; Breton AIDS 2006*

C-IRIS

C-IRIS has been described in people with HIV usually between 2 weeks and 3 months after commencement of ART. Patients develop exaggerated symptoms and signs or atypical inflammation, reminiscent of a paradoxical recurrence,^{126,127} but C-IRIS can also occur in the setting of immune recovery or withdrawal of immunosuppressants. It has also been observed in seemingly immunocompetent individuals, including in *C. gattii* infections, as a post-infectious inflammatory immune response syndrome (PIIRS).^{90,128} There is no diagnostic biomarker for C-IRIS. It is diagnosed by diagnosis of exclusion (figure 2).

There have been no therapeutic trials in C-IRIS. Management strategies include therapeutic lumbar puncture and symptomatic therapies. In severe C-IRIS, corticosteroids are commonly used to dampen inflammation, although their efficacy has not been rigorously examined in clinical trials. In steroid-refractory C-IRIS, there are case reports on the use of tumour necrosis factor- α blockers, such as adalimumab^{129–132} or thalidomide,^{133–135} with mixed success. Corticosteroids can also be beneficial in PIIRS (panel 12; appendix p 76).¹³⁶

Early Adjunctive Steroids ?

Yes

Acute Bacterial Meningitis

De Gans, New Eng J Med 2002

(Reduced mortality and disability)

Tuberculous Meningitis

Thwaites, New Eng J Med 2004

(Reduced mortality)

Donovan, New Eng J Med 2023

(HIV+ patients ?)

Maybe ?

Viral (HSV) Encephalitis

T Solomon, DEXENCEPH

trial
(Improved cognitive function ?)

Listeria monocytogenes meningitis

Brouwer, eClinical Med 2023

(Reduced mortality and disability)

No

Cryptococcal Meningitis

Beardsley, New Eng J Med 2016

(More disability, more adverse events)

Tuberculous meningitis

Drug	WHO recommended daily dose		WHO recommended duration	CSF penetration (CSF/plasma concentration)	Important adverse effects
	Children	Adults			
<i>First-line drugs for the treatment of drug-sensitive TBM</i>					
Rifampicin	15 mg/kg (range 10–20 mg/kg); max. 600 mg	10 mg/kg (range 8–12 mg/kg); max. 600 mg	12 months	10–20%	Hepatotoxicity, orange urine, many drug interactions
Isoniazid	10 mg/kg (range 7–15 mg/kg); max. 300 mg	5 mg/kg (range 4–6 mg/kg); max. 300 mg	12 months	80–90%	Hepatotoxicity, peripheral neuropathy, lupus-like syndrome, confusion, seizures
Pyrazinamide	35 mg/kg (range 30–40 mg/kg)	25 mg/kg (range 20–30 mg/kg)	First 2 months	90–100%	Hepatotoxicity, arthralgia, gout
Ethambutol	20 mg/kg (range 15–25 mg/kg)	15 mg/kg (range 15–20 mg/kg)	First 2 months	20–30%	Dose-related retrobulbar neuritis, more common in renal impairment
Streptomycin*	15–30 mg/kg; max. 1 g IV or IM	15 mg/kg (range 12–18 mg/kg); max. 1 g	First 2 months	10–20%	Monitor plasma concentrations when possible; causes nephrotoxicity and ototoxicity

RJ Wilkinson, Nature Reviews Neurology 2017

Drug	WHO recommended daily dose		WHO recommended duration	CSF penetration (CSF/plasma concentration)	Important adverse effects
	Children	Adults			
<i>Core second-line drugs for the treatment of MDR-TBM[‡]</i>					
Levofloxacin	<5 years: 15–20 mg/kg ≥5 years: 10–15 mg/kg	10–15 mg/kg	Throughout treatment	70–80%	Nausea, headache, tremor, confusion, tendon rupture (rare)
Moxifloxacin	10–20 mg/kg; max. 400 mg Not well established	400 mg	Throughout treatment	70–80%	Nausea, headache, tremor, confusion, tendon rupture (rare)
Amikacin	15–30 mg/kg; max. 1 g IV or IM	15 mg/kg; max. 1 g IV or IM	Intensive phase only	10–20%	Monitor plasma concentrations when possible; causes nephrotoxicity and ototoxicity
Kanamycin	15–30 mg/kg; max. 1 g IV or IM	15 mg/kg; max. 1 g IV or IM	Intensive phase only	10–20%	Monitor plasma concentrations when possible; causes nephrotoxicity and ototoxicity
Capreomycin	15–30 mg/kg; max. 1 g IV or IM	15 mg/kg; max. 1 g IV or IM	Intensive phase only	No data (probably very low)	Monitor plasma concentrations when possible; causes nephrotoxicity and ototoxicity
Ethionamide or prothionamide	15–20 mg/kg; max. 1 g	15–20 mg/kg; max. 1 g	Throughout treatment	80–90%	Anorexia, nausea, vomiting, gynaecomastia, hypothyroidism, confusion, seizures
Cycloserine	10–20 mg/kg; max. 1 g	10–15 mg/kg; max. 1 g	Throughout treatment	80–90%	CNS toxicity: depression, seizures, neuropathy; co-administration with pyridoxine recommended
Linezolid	10 mg/kg; max. 600 mg	600 mg	Throughout treatment	30–70%	Myelosuppression, optic neuropathy; co-administration with pyridoxine recommended

current concepts in anti-tuberculosis treatment in the critically ill patient

Mervyn Mer¹, Alimuddin Zumla² and Martin W. Dünsen^{3*}

Standard anti-TB drugs	
	IV Form
Rifampicin	Yes*
Isoniazid	Yes*
Ethambutol	Yes*
Pyrazinamide	No

Select alternative anti-TB drugs for use in critically ill patients			
	IV Form		IV Form
Levofloxacin	Yes	Linezolid	Yes
Moxifloxacin	Yes	Imipenem + cilastin	Yes
Amikacin	Yes	Meropenem + clavulanate	Yes
Kanamycin	Yes		

*, inconsistent availability; IV, intravenous.

Fig. 1 Potential novel approach to anti-TB treatment in critically ill patients consisting of three (instead of two) phases of therapy. Asterisk indicates inconsistent availability. *IV* intravenous

Adjuvant therapies in critical care: steroids to treat infectious diseases

José Manuel Pereira^{1,2,3*}, Thiago Lisboa^{4,5} and José-Artur Paiva^{1,2,3}

© 2017 Springer-Verlag GmbH Germany, part of Springer Nature and ESICM

In tuberculosis

A review of 1337 patients with mild to severe tuberculous meningitis showed a quarter mortality reduction with steroids [RR 0.75 (CI 0.65 to 0.87)] [21], at least in the short term. The survival benefit was mostly seen in patients with mild disease and CS have little or no effect if advanced neurologic symptoms are present. In addition, no increased risk of adverse effects was observed. Therefore, adjunctive CS, such as dexamethasone tapering over 4 weeks from 0.3 to 0.4 mg/kg, is recommended for tuberculous meningitis.