Best-of

Infections sur prothèses vasculaires

Dr Simon JAMARD

CHRU de Tours – Hôpital Bretonneau

Service de Médecine Infectieuse et Tropicale

INRAE – UMR 1282 – Infectiologie et Santé Publique

Déclaration d'intérêt de 2014 à 2025

- Intérêts financiers : aucun
- Liens durables ou permanents : aucun
- Interventions ponctuelles : orateur atelier Philips
- Intérêts indirects : aucun

Actualité sur les IPV / Perplexity

quelles sont les dernière nouveauté dans la littérature scientifique pour la prise en charge des infections sur prothèses vasculaires

Réponse

Images

Sources

Tâches

Vascular Graft Infection Following Aneurysm Repair: A Population-Based Study

Hussam Tabaja, M.D.¹, Larry M. Baddour, M.D.^{1,2}, Supavit Chesdachai, M.D.¹, Randall R. DeMartino, M.D.³, Brian D. Lahr, M.S.⁴, Daniel C. DeSimone, M.D.^{1,2}

2023

- ❖ Mayo Clinic – 2010 à 2020
- ❖ 708 interventions
 - 291 Chirurgies ouvertes
 - 417 Endovasculaires
- ❖ Suivi médian : 4,1 [1,9-6,8] années
- ❖ Infections : 15 épisodes
- ❖ Incidence cumulative à 5 ans : 1,6% [0,6-2,7%]
 - Endovasculaire : 1,4% [0,2-2,6%]
 - Chirurgie ouverte : 2,0% [0,3-3,7%]

Caractéristiques des IPV

Characteristics and Outcome of Vascular Graft Infections: *Open Forum Infectious Diseases* A Risk Factor and Survival Analysis

MAJOR ARTICLE

2024

Leonie Stockschläder,^{1,*} Donara Margaryan,¹ Safwan Omran,^{2,*} Martin Schomaker,² Andreas Greiner,² and Andrei Tramnuz^{1,*}

❖ Cohorte
rétrospective

❖ 2010-2020

❖ Berlin

Characteristic	All Patients (n = 78)	Aortic VGI (n = 49)	Peripheral VGI (n = 29)	P Value
Onset of infection				
Early-onset infection (<8 wk)	30 (38.5)	15 (30.6)	15 (51.7)	.064
Late-onset infection (≥8 wk)	48 (61.5)	34 (69.4)	14 (48.3)	.064
Time from initial graft implantation to infection diagnosis, d, median (range)	160 (1–7928)	363 (6–7928)	56 (1–2638)	.018
Clinical signs and symptoms at the time of initial diagnosis				
Fever	19 (24.4)	18 (36.7)	1 (3.4)	.001
Fatigue	7 (9.0)	7 (14.3)	0 (0.0)	.033
Surgical site erythema	23 (29.5)	8 (16.3)	15 (51.7)	.001
Surgical site discharge	22 (28.2)	12 (24.5)	10 (34.5)	.343
Local pain	22 (28.2)	11 (22.4)	11 (37.9)	.142
Graft thrombosis/stenosis	16 (20.5)	5 (10.2)	11 (37.9)	.003
Radiologic findings at the time of initial diagnosis				
Perigraft fluid	22 (28.2)	21 (42.9)	1 (3.4)	<.001
Perigraft air	15 (19.2)	12 (24.5)	3 (10.3)	.126
Perigraft abscess	15 (19.2)	11 (22.4)	4 (13.8)	.349
Graft thrombosis/stenosis	14 (17.9)	5 (10.2)	9 (31.0)	.021
Perigraft enhancement in PET/CT	12 (15.4)	12 (24.5)	0.0 (0)	.004
Aneurysm involving or surrounding the graft	10 (12.8)	6 (12.2)	13.8 (4)	.843
Perigraft lymphadenopathy	9 (11.5)	9 (18.4)	0 (0.0)	.014
Perigraft hematoma	7 (9.0)	6 (12.2)	1 (3.4)	.189
Anastomotic insufficiency/dehiscence/rupture	4 (5.1)	4 (8.2)	0 (0.0)	.291
Graft-organ fistula	2 (2.6)	2 (4.1)	0 (0.0)	.527

Caractéristiques des IPV

Characteristics and Outcome of Vascular Graft Infections: *Open Forum Infectious Diseases* A Risk Factor and Survival Analysis

MAJOR ARTICLE

2024

Leonie Stockschläder,^{1,*} Donara Margaryan,¹ Safwan Omran,^{2,*} Martin Schomaker,² Andreas Greiner,² and Andrej Trampuz^{1,*}

- ❖ Cohorte rétrospective
- ❖ 2010-2020
- ❖ Berlin

Pathogen	All Patients (n = 78)	Aortic VGI (n = 49)	Peripheral VGI (n = 29)	P Value
<i>Staphylococcus</i> spp	30 (38.5)	15 (31.3)	15 (51.7)	.074
Coagulase-negative staphylococci ^a	14 (18.0)	7 (14.3)	7 (24.1)	.273
<i>S aureus</i>	15 (19.2)	7 (14.3)	8 (27.6)	.163
Gram-negative bacteria ^b	25 (32.1)	13 (27.1)	12 (41.1)	.194
<i>Enterococcus</i> spp	20 (25.6)	11 (22.9)	9 (31.0)	.431
<i>Candida</i> spp ^c	17 (21.8)	12 (25.0)	5 (17.2)	.426
Anaerobes ^d	8 (10.3)	6 (12.2)	2 (6.9)	.452
Streptococci	7 (9.0)	3 (6.3)	4 (13.8)	.265
<i>Pseudomonas</i> spp	7 (9.0)	2 (4.2)	5 (17.2)	.053
Drug-resistant pathogens ^e	7 (9.0)	5 (10.4)	2 (6.9)	.603
Polymicrobial	32 (41.6)	19 (39.6)	13 (44.8)	.651
Culture negative	14 (18.2)	13 (27.1)	1 (3.4)	.009

Caractéristiques des IPV

Characteristics and Outcome of Vascular Graft Infections: A Risk Factor and Survival Analysis

MAJOR ARTICLE

2024

Leonie Stockschläder,^{1,✉} Donara Margaryan,¹ Safwan Omran,^{2,✉} Martin Schomaker,² Andreas Greiner,² and Andrej Trampuz^{1,✉}

IPV fongiques

Fungal versus non-fungal supra-inguinal prosthetic vascular graft infections: A cohort study

Baptiste Monnier ^a, Thibault Couture ^b, Agnès Dechartres ^c, Samuel Sitruk ^d, Johann Gaillard ^e, Alexandre Bleibtreu ^f, Laurent Chiche ^g, Julien Gaudric ^b, Jérémy Arzoine ^h

2024

- ❖ Cohorte retrospective 2009-2021
- ❖ 32 IPV fongiques vs 116 IPV non fongiques

Variable	Multivariable		
	OR	95% IC	p value
Prostheto-digestive fistula	5.98	(2.29-15.62)	< 0.001
Preoperative antibiotic therapy ≥ 7 days	2.87	(1.12-7.38)	0.03
Immunocompromised	1.45	(0.39-5.38)	0.58
Late infection > 4 months	2.54	(0.86-7.52)	0.09
Preoperative use of norepinephrine	3.98	(0.95-16.61)	0.06

Abbreviations: OR, odds ratio; CI, confidence interval.

Critères diagnostics

Editor's Choice – Validation of the Management of Aortic Graft Infection Collaboration (MAGIC) Criteria for the Diagnosis of Vascular Graft/Endograft Infection: Results from the Prospective Vascular Graft Cohort Study

2021 Alexia Anagnostopoulos ^{a,*,†}, Fabienne Mayer ^{a,†}, Bruno Ledergerber ^a, Judith Bergada-Pijuan ^a, Lars Husmann ^b, Carlos A. Mestres ^c, Zoran Rancic ^d, Barbara Hasse ^{a,*}, the VASGRA Cohort Study

Table 5. The MAGIC classification¹

Criterion	Clinical/surgical	Radiology	Laboratory
<i>Major</i>	Pus (confirmed by microscopy) around graft or in aneurysm sac at surgery	Perigraft fluid on CT scan $\geq$ 3 months after insertion	Organisms recovered from an explanted graft
	Open wound with exposed graft or communicating sinus	Perigraft gas on CT scan $\geq$ 7 weeks after insertion	Organisms recovered from an intra-operative specimen
	Fistula development, e.g., aorto-enteric or aortobronchial	Increase in perigraft gas volume demonstrated on serial imaging	Organisms recovered from a percutaneous, radiologically guided aspirate of perigraft fluid
<i>Minor</i>	Graft insertion in an infected site, e.g., fistula, mycotic aneurysm, or infected pseudo-aneurysm	Other, e.g., suspicious perigraft gas/fluid soft tissue inflammation; aneurysm expansion; pseudo-aneurysm formation: focal bowel wall thickening; discitis/osteomyelitis; suspicious metabolic activity on FDG-PET/CT; radiolabelled leukocyte uptake	Blood culture(s) positive and no apparent source except graft infection
	Localised clinical features of graft infection, e.g., erythema, warmth, swelling, purulent discharge, pain		
	Fever $\geq$ 38°C with graft infection as most likely cause		Abnormally elevated inflammatory markers with graft infection as most likely cause, e.g., erythrocyte sedimentation rate, C reactive protein, white cell count

❖ Validation score Magic sur cohorte existante VASGRA

❖ 137 patients avec IPV
120 contrôles

Critères diagnostics

Editor's Choice – Validation of the Management of Aortic Graft Infection Collaboration (MAGIC) Criteria for the Diagnosis of Vascular Graft/Endograft Infection: Results from the Prospective Vascular Graft Cohort Study

2021 Alexia Anagnostopoulos ^{a,*,†}, Fabienne Mayer ^{a,†}, Bruno Ledergerber ^a, Judith Bergada-Pijuan ^a, Lars Husmann ^b, Carlos A. Mestres ^c, Zoran Rancic ^d, Barbara Hasse ^{a,*}, the VASGRA Cohort Study

MAGIC adjudication	VASGRA		Sensitivity (95% CI) – %	Specificity (95% CI) – %		Total
	Confirmed	<i>“Diseased” = definite and suspected VGEI; “not diseased” = rejected VGEI and controls</i>			Control patients	
Confirmed VGEI	126 (93.3)	Overall	99 (96–100)	61 (52–70)	3 (3)	135 (52.5)
Suspected VGEI	8 (5.9)	Intracavitary abdominal VGEI	100 (93–100)	62 (51–72)	14 (16)	48 (18.7)
Excluded VGEI	1 (0.7)	Thoracic aorta VGEI	98 (88–100)	42 (15–72)	0 (0)	6 (2.3)
Control patients	0 (0)	Peripheral arteries VGEI	NA	N/A	68 (80)	68 (26.4)
Total	135 (100)	<i>“Diseased” = definite VGEI; “not diseased” = suspected VGEI, rejected VGEI, and controls</i>			85 (100)	257 (100)
		Overall	93 (88–97)	93 (87–97)		
		Intracavitary abdominal VGEI	94 (84–99)	92 (85–96)		
		Thoracic aorta VGEI	86 (73–95)	100 (74–100)		
		Peripheral arteries VGEI	100 (91–100)	67 (9.0–99)		

Diagnostic microbiologique

Sonication of Vascular Grafts and Endografts to Diagnose Vascular Graft Infection: a Head-To-Head Comparison with Conventional Culture and Its Clinical Impact

Lisanne Braams,^a Gro Vlaspolder,^a Kathleen Boiten,^a Elisa Salomon,^a Rik Winter,^a Ben Saleem,^b Marjan Wouthuyzen-Bakker,^a Marleen van Oosten^a

TABLE 2 Diagnostic performance^c

Method	Episodes (N = 40)		Grafts (N = 57)		Cultured causative microorganisms (N = 107) and contaminants (N = 26)	
	Positive results/VGEI ^a	Negative results/non-VGEI ^a	Positive results/VGEI ^a	Negative results/non-VGEI ^a	Causative m.o./total causative m.o.	Contamination rate (contaminants/total m.o. per method)
Conventional culture	26/32	5/8	34/44	8/13	90/107	15/105 (14%)
Sonication culture	26/32 ^b	7/8 ^b	33/44 ^b	13/13 ^b	71/107 ^b	11/82 (13%) ^b

❖ 32 % de Mismatch

- Anaérobie plus souvent retrouvé en culture conventionnelle
- *E. faecalis*, *E. corrodens*, Lactobacilli retrouvé sur sonicat

Diagnostic microbiologique

Sonicate Fluid Inoculation Into Blood Culture Bottles for Vascular Graft and Endograft Infection Microbiological Diagnosis

Mathilde Puges ^{a,c*}, Mathilde Carrer ^a, Caroline Caradu ^b, Xavier Bérard ^b, Charles Cazanave ^{a,c}, on behalf of the VIP study group

2024

❖ Inoculation du sonicat dans un flacon d'hémoculture

Variable	Sonicate fluid culture	SF-BCBs	Combined
Microbiological diagnosis	24 (64.9)	24 (64.9)	26 (70.3)
Isolated microorganisms in total (excluding potential contaminants)	47	34	
Patients with polymicrobial infections	11 (29.7)	7 (18.9)	12 (32.4)
Isolated pathogenic microorganisms per patient	2.08 (1–7)	1.48 (1–4)	2.08 (1–7)
Isolated microorganisms			
Enterobacteriaceae	13	12	
Staphylococci	13	10	
<i>Staphylococcus aureus</i>	6	5	
Coagulase negative staphylococci	7	5	
Streptococci	5	4	
Anaerobes	5	2	
Enterococci	4	3	
Non fermenting Gram negative bacilli	3	2	
<i>Candida</i> spp.	2	1	
<i>Lactobacillus</i> spp.	1	0	
<i>Corynebacterium amycolatum</i>	1	0	
Patients with supplementary microorganisms identified	8 (21.6)	4 (10.8)	
Modification of anti-infectious treatment for supplementary microorganisms	4 (14.8)	3 (8.1)	
Median delay of first microbial isolation – hours	27 [15.6–45]	10 [4.7–19]	12 [5–19]
Median delay of first microbial identification – hours	31 [18.5–48]	21.3 [18.5–30]	20.5 [17.3–24.8]
Patients with a quickest microbial identification	9 (45)	11 (55)	
Patients with > 1 potential contaminant microorganisms*	9 [†] (24.3)	9 [†] (24.3)	16 (43.2)

Biofilm et IPV

2023

Biofilm formation on endovascular aneurysm repair (EVAR) grafts—a proof of concept *in vitro* model

Torgny Sunnerhagen^{1,2,3} ✉, Franziska Schwartz¹, Lars Christophersen¹,
Thomas Bjarnsholt^{1,4}, Klaus Qvortrup⁵, Nikolaj Eldrup⁶, Katja Vogt⁶, Claus Moser^{1,4}

Biofilm et IPV

Efficacy of rifampicin combination therapy against MRSA prosthetic vascular graft infections in a rat model

Mikkel Illemann Johansen ^{a,b,1}, Maiken Engelbrecht Petersen ^{c,1}, Emma Faddy ^{a,b}, Anders Marthinsen Seefeldt ^b, Alexander Alexandrovich Mitkin ^c, Lars Østergaard ^{b,a}, Rikke Louise Meyer ^{c,d,*}, Nis Pedersen Jørgensen ^{b,**}

2024

Prise en soin

Merits of Puncture and Drainage of Perigraft Material in Patients with Vascular and Endovascular Prosthetic Graft Infection

In press [Giulia Colonna](#)^a, [Reinoud P.H. Bokkers](#)^b, [Marjan Wouthuyzen Bakker](#)^c, [Barzi Gareb](#)^d, [Maurizio Taurino](#)^a, [Clark J. Zeebregts](#)^e, [Ben R. Saleem](#)^e

- ❖ Cohorte rétrospective – 2018-2023 – Pays bas
- ❖ 69 patients inclus
 - 45 IPV abdominale
 - 24 IPV périphériques
- ❖ Score ASA ≥ 3
- ❖ Aspiration percutanée et/ou mise en place d'un drain dans la collection périprothétique
- ❖ Complications :
 - Drainage : 2 plaies rénales et 1 décès sur choc septique

Table 6. Microbiological mismatch between micro-organisms detected in percutaneous cultures or blood cultures vs. micro-organisms detected in extracted vascular grafts from patients with vascular graft or endograft infections (VGEIs).

Match/mismatch	Abdominal VGEI	Peripheral VGEI
<i>Percutaneous cultures vs. vascular graft</i>		
Total	21	11
<i>Mismatch</i>	10 (48)	8 (73)
Different micro-organism(s)	2 (20)	4 (50)
Percutaneous culture negative	5 (50)	4 (50)
Vascular graft culture negative	3 (30)	0 (0)
<i>Match</i>	11 (52)	3 (27)
Same micro-organism(s)	2 (18)	0 (0)
Partially same micro-organisms	7 (64)	3 (100)
Both cultures negative	2 (18)	0 (0)
<i>Blood cultures vs. vascular graft</i>		
Total	12	7
<i>Mismatch</i>	7 (58)	7 (100)
Different micro-organism(s)	0 (0)	0 (0)
Blood culture negative	6 (86)	7 (100)
Vascular graft culture negative	1 (14)	0 (0)
<i>Match</i>	5 (42)	0 (0)
Same micro-organism(s)	1 (20)	0 (0)
Partially same micro-organisms	2 (40)	0 (0)
Both cultures negative	2 (40)	0 (0)

Data are presented as n (%). VGEI = vascular graft or endograft infection.

Prise en soin

Merits of Puncture and Drainage of Perigraft Material in Patients with Vascular and Endovascular Prosthetic Graft Infection

In press

[Giulia Colonna](#)^a , [Reinoud P.H. Bokkers](#)^b, [Marjan Wouthuyzen Bakker](#)^c, [Barzi Gareb](#)^d, [Maurizio Taurino](#)^a, [Clark J. Zeebregts](#)^e, [Ben R. Saleem](#)^e

- ❖ Cohorte rétrospective – 2018-2023 – Pays bas
- ❖ 69 patients inclus
 - 45 IPV abdominale
 - 24 IPV périphériques
- ❖ Score ASA ≥ 3
- ❖ Aspiration percutanée et/ou mise en place d'un drain dans la collection périprothétique
- ❖ Complications :
 - Drainage : 2 plaies rénales et 1 décès sur choc septique

Prise en soin

Semi-Conservative Treatment Versus Radical Surgery in Abdominal Aortic Graft and Endograft Infections

Hamid Gavali ^{a,*}, Kevin Mani ^a, Mia Furebring ^b, Karl Wilhelm Olsson ^a, David Lindström ^a, Karl Sörelius ^c, Birgitta Sigvant ^{a,d}, Gustav Torstensson ^e, Manne Andersson ^{f,g}, Claes Forssell ^h, Håkan Åstrand ^g, Tobias Lundström ⁱ, Shahzad Khan ^j, Björn Sonesson ^j, Otto Stackelberg ^{k,l}, Peter Gillgren ^{k,m}, Jon Isaksson ⁿ, Björn Kragsterman ^{a,o}, Khatereh Djavani Gidlund ^p, Tal Horer ^p, Mitra Sadeghi ^p, Anders Wanhainen ^{a,n}

2023

Baseline characteristic	SC (n = 43)	RS (n = 126)	p value
Median age (IQR) – y	73.6 (67.1, 78.4)	70.0 (65.3, 75.7)	.039
Male sex	35 (81.4)	104 (82.5)	.98
Hypertension	28 (65.1)	79 (65.3)	1.0
Smoking	7 (16.3)	48 (40.3)	.005
Chronic kidney disease	9 (20.9)	13 (10.6)	.092
Diabetes	8 (18.6)	10 (8.1)	.062
Heart failure	10 (23.3)	9 (7.4)	.010
Coronary artery disease	26 (60.5)	37 (30.6)	<.001
Lung disease	8 (18.6)	13 (10.7)	.19
Circulatory shock	7 (16.3)	11 (8.7)	.19
ASA score >3 *	22 (53.7)	33 (27.3)	.002
Graft enteric fistula	23 (53.5)	62 (50.4)	.73

Complication	SC (n = 43)	RS (n = 126)	p value
Median follow up (IQR) – y [±]	5.1 (3.9, 14.6)	3.9 (3.2, 9.5)	.18
Recurrent graft infection	45.4	19.3	<.001
Anastomosis dehiscence or stump blowout	2.3	10.5	.10
Amputation	9.3	4.3	.23
Re-intervention	23.3	26.3	.70

No. at risk						
— Semi-conservative	43	25	21	18	15	13
— Radical surgery	126	87	81	72	65	60

Prise en soin

Biodesign: Engineering an aortic endograft explantation tool

Solyman Hatami, MD,^a Vamsi Maturi, MD,^a Alwin Mathew, MD,^a Shannon Lu, MD,^a Paul Haddad, MD,^b Daanish Sheikh, BS,^c and Maham Rahimi, MD, PhD,^b Houston and San Antonio, TX

2024

2024 Fostering Collaborative Teamwork—A Comprehensive Approach to Vascular Graft Infection Following Arterial Reconstructive Surgery

Hussam Tabaja,^{1,6} Supavit Chesdachai,^{1,6} Aditya S. Shah,^{1,6} Ryan W. Stevens,² Randall R. DeMartino,³ Young M. Er Larry M. Baddour,^{1,5,6} and Daniel C. DeSimone^{1,5}

26^{es} JNi 2025
TOURS

et la région Centre - Val de Loire VINCI

Journées Nationales d'infectiologie

du mercredi 11 juin 2025 au vendredi 13 juin 2025

Journée Nationale de Formation
des Paramédicaux en Infectiologie

Jeudi 12 juin 2025

Merci pour votre attention !

Séminaire du

Groupe
Infectiologie
Digit@le

Projets et RETEX

Ateliers pratiques

Table ronde Implémentation

Best-of digital

Octobre 2025

Save the date!